Fluid Phase Equilibria, 95 (1994) 401–402
Elsevier Science B.V.

Contents of Volume 95 (1994)

Equation of state calculations and Monte Carlo simulations of internal energies, compressibility factors and vapor–liquid equilibria for Lennard-Jones fluid mixtures
Y. Miyano (Okayama, Japan) 1–13

Solid–solid transition and solubility of even *n*-alkanes
A. Książczak, K. Moorthi and I. Nagata (Kanazawa, Japan) 15–29

An effective triplet potential for argon
Y. Miyano (Okayama, Japan) 31–41

A model for predicting the excess enthalpies of ternary mixtures
L. Wang, S. Shen, G.C. Benson and B.C.-Y. Lu (Ottawa, Ont., Canada) 43–57

On the relationship between the equilibrium constants of consecutive association reactions
A. Anderko and J.M. Prausnitz (Berkeley, CA, USA) 59–71

A model-free method for calculating vapor–liquid equilibria for multicomponent systems from total-pressure or boiling-point data
Y. Hu, H. Liu (Shanghai, China) and J.M. Prausnitz (Berkeley, CA, USA) 73–92

Non-iterative flash calculation algorithm in compositional reservoir simulation
P. Wang and E.H. Stenby (Lyngby, Denmark) 93–108

A partial molar fugacity coefficient useful for changing mixing rules of cubic equations of state
A.S. Lawal and E.T. van der Laan (Anchorage, AK, USA) 109–121

VLE data and UNIFAC-parameters of binary mixtures with methylisocyanate
P. Ackermann, K.F. Knoche and M. Roth (Aachen, Germany) ... 123–136

Solubility of CCl_2F_2, $CClF_3$, CF_4 and c-C_4F_8 in H_2O and D_2O at 288 to 318 K and 101.325 kPa. Thermodynamics of transfer of gases from H_2O to D_2O
P. Scharlin (Turku, Finland) and R. Battino (Dayton, OH, USA) . 137–147

Infinite-dilution activity coefficients by comparative ebulliometry. Binary systems containing chloroform and diethylamine
J.P. Coutinho and E.A. Macedo (Porto, Portugal) 149–162

Gas solubility of hydrogen sulfide and carbon dioxide in mixtures of sulfolane with diethanolamine at different temperatures
F. Murrieta-Guevara, E. Rebolledo-Libreros and A. Trejo (Mexico, Mexico) 163–174

Application of the UNIFAC and Nitta–Chao models to describing the behavior of methyl ester/alkane mixtures, and experimental data for (methyl *n*-alkanoates + *n*-heptadecane) binary mixtures
J. Ortega (Las Palmas de Gran Canaria, Spain) and J.L. Legido (Vigo, Spain) 175–214

Solubilities of the xanthines caffeine, theophylline and theobromine in supercritical carbon dioxide
M. Johannsen and G. Brunner (Hamburg, Germany) 215–226

(p, V_m, T, x) measurements for liquid mixtures of 1,2-dichloroethane with 2,2,4-trimethylpentane. I. Experimental results, isothermal compressibilities, isobaric expansivities and heat capacities
R. Malhotra and L.A. Woolf (Canberra, A.C.T., Australia)..... 227–251
(p, V_m, T, x) measurements for liquid mixtures of 1,2-dichloroethane with 2,2,4-trimethylpentane. II. Thermodynamic excess properties
R. Malhotra and L.A. Woolf (Canberra, A.C.T., Australia)..... 253–266
Solubility and volumetric behavior of methane and carbon dioxide plus C_{7+} pseudocomponents of an oil
S.-P. Hong, K.A. Green and K.D. Luks (Tulsa, OK, USA)..... 267–280
The water content of ethane, propane and their mixtures in equilibrium with liquid water or hydrates
K.Y. Song and R. Kobayashi (Houston, TX, USA)..... 281–298
Excess molar volumes and viscosities of binary mixtures of bis(2-methoxyl)ether (diglyme) with chloroalkanes at 298.15 K
M.E. de Ruiz Holgado, C.R. de Schaefer, E.L. Arancibia and M. Katz (Tucumán, Argentina)..... 299–312
High pressure phase equilibria of the system phenol–*m*-cresol–carbon dioxide
G. Di Giacomo, S. Brandani, V. Brandani and G. Del Re (L'Aquila, Italy)..... 313–327
Virial coefficients of *n*-butane from pressure measurement only
B.D. Ababio, P.J. McElroy and A.G. Williamson (Christchurch, New Zealand)..... 329–339
Thermodynamics of potassium hydrogen phosphate–potassium dihydrogen phosphate–polyethylene glycol aqueous two-phase systems
Q. Peng, Z. Li and Y. Li (Beijing, China)..... 341–357

Short Communications

Solid–liquid–vapour equilibrium of binary systems phenol + hexane, phenol + heptane and 4-methylphenol + hexane
A. Książczak (Warsaw, Poland) and I. Nagata (Kanazawa, Japan) 359–369
Influence of molecular shape on vapour–liquid equilibria. Application of a modified Christoforakos–Franck equation of state
Đ. Plačkov, D.E. Mainwaring and R.J. Sadus (Hawthorn, Vic., Australia)..... 371–381

Review

Prediction of parachors of petroleum cuts and pseudocomponents
J.K. Ali (Loughborough, UK)..... 383–398

Subject Index..... 399

Author Index..... 400

Contents of Volume 95 (1994)..... 401–402